

# Hemothorax Management After Endovascular Treatment For Thoracic Aortic Rupture

G. Piffaretti <sup>a,\*</sup>, M. Menegolo <sup>b</sup>, A. Kahlberg <sup>c</sup>, G. Mariscalco <sup>d</sup>, E. Rinaldi <sup>c</sup>, P. Castelli <sup>a</sup>, F. Grego <sup>b</sup>, R. Chiesa <sup>c</sup>, M. Antonello <sup>b</sup>

<sup>a</sup> Department of Surgery and Morphological Sciences, University of Insubria School of Medicine, Circolo University Hospital, Varese, Italy

<sup>b</sup> Department of Cardiac, Thoracic and Vascular Sciences, University of Padua School of Medicine, Padua University Hospital, Padua, Italy

<sup>c</sup> Department of Vascular Surgery, "Vita Salute" University School of Medicine, San Raffaele Scientific Institute, Milan, Italy

<sup>d</sup> Cardiac Surgery, Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Leicester, United Kingdom

## WHAT THIS PAPER ADDS

This study has focused on the characteristics and management of hemothorax, occurring in a large cohort of patients with thoracic aortic rupture, treated with TEVAR. It may contribute further insights into the management of ruptured descending thoracic aorta and into optimizing the post-operative risk stratification after urgent TEVAR. Furthermore, it is hoped it will lead to larger studies to optimize current practice on the best management of hemothorax after TEVAR.

**Objectives:** The aim was to describe and analyze the management of hemothorax (HTX) and the occurrence of respiratory complications after endovascular repair of thoracic aortic rupture (TEVAR).

**Methods:** This was a multicenter study with retrospective analysis. Between November 2000 and December 2012, all patients with confirmed HTX due to rupture of the descending thoracic aorta treated with TEVAR were included. Respiratory function (acid base status, PaO<sub>2</sub>, Paco<sub>2</sub>, lactate, and respiratory index) was monitored throughout hospitalization. Primary endpoints were survival and post-operative respiratory complications.

**Results:** Fifty-six patients were treated. The mean age was 62 ± 21 years (range 18–92 years). Etiology included traumatic rupture (*n* = 23, 41%), atherosclerotic aneurysm (*n* = 20, 36%), Debakey type IIIa dissection (*n* = 8, 14%), and penetrating aortic ulcer (*n* = 5, 9%). The primary technical success of TEVAR was 100%. The in hospital mortality rate was 12.5% (*n* = 7). Hemothorax was drained in 21 (37.5%) cases. In hospital respiratory complications occurred in 23 (41%) patients who required a longer intensive care unit stay (days 2.3 ± 0.7 vs. 1.9 ± 0.8, *p* = .017), and hospitalization (26 ± 17 vs. 19 ± 17, *p* = .021). Those who developed post-operative respiratory complications had lower pre-operative PO<sub>2</sub> values (mmHg, 80 ± 24 vs. 91 ± 21, *p* = .012). Respiratory complications and in hospital mortality did not differ among aortic pathologies (*p* = .269 and *p* = 1.0, respectively), nor did in hospital mortality differ between patients with and without respiratory complications (13% vs. 12%; *p* = .990).

**Conclusions:** Thoracic aortic rupture still has a high mortality rate. Respiratory complications have not been eliminated by endovascular repair. HTX evacuation may have had a positive influence on the survival in these patients. Although traumatic and degenerative ruptures are two significantly different scenarios, survival and respiratory outcomes were similar and were not affected by the underlying aortic disease.

© 2015 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

Article history: Received 4 September 2014, Accepted 8 July 2015, Available online 8 September 2015

**Keywords:** Hemothorax, TEVAR, Respiratory complications, Thoracic aortic endovascular repair, Thoracic aortic rupture

## INTRODUCTION

Rupture of the descending thoracic aorta has traditionally been treated by open graft replacement, with its associated mortality and complication rates which remain disturbingly high.<sup>1–3</sup> More recently, thoracic endovascular aortic repair (TEVAR) has proved to be an effective, less traumatic, alternative for the treatment of all thoracic aortic pathologies; indeed several papers have advocated TEVAR as the first choice for the emergency treatment of thoracic aortic catastrophes.<sup>4–7</sup> Rupture of the thoracic aorta is frequently associated with hemothorax (HTX); however, the

DOI of original article: <http://dx.doi.org/10.1016/j.ejvs.2015.08.001>

\* Corresponding author. Department of Surgery and Morphological Sciences, Circolo University Hospital, University of Insubria School of Medicine, Via Guicciardini, 9, 21100 Varese, Italy.

E-mail address: [gabriele.piffaretti@uninsubria.it](mailto:gabriele.piffaretti@uninsubria.it) (G. Piffaretti).

1078-5884/© 2015 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.ejvs.2015.07.039>

management of HTX after TEVAR has not been debated until now, and there is still uncertainty on indication and timing of its treatment.<sup>8–14</sup> Hemothorax may lead to major complications such as compression of the esophagus and/or cardiovascular structures, respiratory insufficiency, and infection. All these complications may compromise the post-operative survival in an already critically ill subset of patients.<sup>8</sup> The purpose of this study was to analyze the management of HTX in patients with descending thoracic aortic rupture treated with TEVAR.

## MATERIALS AND METHODS

### Study population

The study was designed as a multicenter retrospective study. Between November 2000 and December 2012, all patients presenting with rupture of the descending thoracic aorta and HTX treated with TEVAR were identified and included in the analysis. Thoraco-abdominal lesions were excluded. Clinical and procedural data were collected prospectively and recorded in a dedicated database. All patients underwent pre-operative thoraco-abdominal computed tomography (CT) scans. Intervention was performed in the operating theatre, equipped to perform either open surgical or endovascular procedures. General anesthesia and antibiotic prophylaxis with a second generation cephalosporin was used in all patients. Four different thoracic endografts (EGs) were implanted: Excluder/TAG/C-TAG (W.L. Gore and Associates, Flagstaff, AZ, USA), Talent/Valiant/Captivia (Medtronic Vascular, Santa Rosa, CA, USA), TX-1/TX-2 (Cook, Bloomington, IN, USA), and Relay (Bolton Medical, Sunrise, FL, USA). Post-operatively, the patient was transferred to the intensive care unit (ICU). In all patients, respiratory function was monitored at admission and during hospitalization: acid base status,  $P_{aO_2}$ ,  $P_{aCO_2}$ , lactate, and the respiratory index ( $P_{aO_2}/F_{iO_2}$ ) were evaluated.<sup>15</sup> Evacuation of the HTX was always discussed case by case by the operating team and the intensive care unit (ICU) physicians; generally, HTX was drained when the respiratory index (RI) was  $< 200$  or when signs of respiratory or cardiovascular impairment were detected. The hemothorax was drained with a chest tube; video assisted thoracoscopy or open thoracotomy which was used selectively for complicated trapped HTX. Clinical and CT scan follow up were performed at 1, 6, and 12 months after the intervention, and annually thereafter.

### Definition

Thoracic aortic rupture was defined as hemorrhage outside the boundaries of the aorta. Hemothorax was considered as any collection of blood in the pleural cavity, but isolated peri-aortic/mediastinal hematoma suggestive for a contained rupture was not counted as HTX. Comorbidities were defined according to the Society of Thoracic Surgeons Adult Database definitions.<sup>16</sup> On admission, hemorrhagic shock was defined as a combination of systolic blood pressure  $< 80$  mmHg after fluid resuscitation or need for  $\alpha$ -amines, tachycardia ( $> 110$  bpm), anuria or urine output  $< 15$  mL/

hour, unconsciousness, or circulatory arrest.<sup>17</sup> The HTX at its point of greater thickness in the baseline CT was classified:  $< 2$  cm, 2–3 cm,  $> 3$  cm. Delay was defined as the time interval between the diagnosis of aortic rupture/onset of symptoms and the start of the intervention. Traumatic aortic injury severity grading was classified accordingly to the clinical practice guidelines of the Society for Vascular Surgery (SVS).<sup>18</sup> The operative risk profile was estimated according to the EuroSCORE.<sup>19</sup> Operative outcomes were classified following the ad hoc committee on TEVAR reporting standards of the Society for Vascular Surgery/American Association for Vascular Surgery (SVS/AAVS).<sup>20</sup> Specifically, respiratory complications were defined as grade 1 if recovery was prompt with medical treatment, grade 2 for prolonged hospitalization or intravenous antibiotics, and grade 3 for prolonged intubation, tracheotomy, deterioration in pulmonary function,  $O_2$  dependence, or fatal outcome. Primary endpoints were the evaluation of survival and respiratory complications.

### Data analysis

Clinical data were recorded in Microsoft Excel (Microsoft Corp, Redmond, WA, USA). Results are presented as mean  $\pm$  SD for continuous variables, and number (percentage) for categorical variables. Continuous variables were compared with the Mann–Whitney U test, and categorical variables by the Fisher exact test. The survival rate was estimated by means of the Kaplan–Meier method with the log-rank test. A  $p$  value  $< .05$  was considered significant. Statistical analysis was computed with SPSS, release 20.0, for Windows (IBM SPSS Inc., Chicago, IL, USA).

## RESULTS

### General data

During the study period a total of 500 TEVARs for descending thoracic aortic diseases were performed: 402 (81%) intact aortic lesions were treated; 42 (8%) were ruptured or traumatic lesions without HTX. Rupture with HTX was present in 56 (11%) cases: there were 40 (71%) males. Mean age was  $62 \pm 21$  years (range 18–92 years). Traumatic ruptures ( $n = 23$ , 41%) were caused by blunt injuries in all cases; a grade IV injury was present in all cases. In particular, traumatic rupture involved the distal aortic arch below the left subclavian artery ( $n = 9$ , 39%) and the descending aorta ( $n = 14$ , 61%). Non-traumatic ruptures were caused by atherosclerotic aneurysm ( $n = 20$ , 36%), Debakey type IIIa dissection ( $n = 8$ , 14%), and penetrating aortic ulcer ( $n = 5$ , 9%). Overall demographic data, co-morbidities and risk factors are reported in Table 1.

### Operative data

Emergent TEVAR was performed in 19 (34%) cases: all but three (5%) patients were treated within 3 hours of rupture with a median delay of 1.5 hours (range 0.5–24). The proximal landing zone was “zone 2” in 18 (32%) cases and “zones 3 or 4” in 38 (68%). Primary technical and primary

**Table 1.** Demographic and presentation data.

| Variable <sup>a</sup>                 | All patients    |
|---------------------------------------|-----------------|
| Demographic data                      |                 |
| M : F                                 | 40 : 16         |
| Age (mean, years)                     | 62 ± 21         |
| Risk factors (%)                      |                 |
| CAD                                   | 16 (29)         |
| COPD                                  | 16 (27)         |
| Obesity (BMI > 30)                    | 15 (27)         |
| Diabetes                              | 13 (23)         |
| AKI                                   | 8 (14)          |
| Valve disease/LHF                     | 7 (13)          |
| Operative risk                        |                 |
| Shock (%)                             | 19 (34)         |
| EuroSCORE (mean, predicted mortality) | 9 ± 4 (35 ± 23) |

AKI = acute kidney injury; BMI = body mass index; CAD = coronary artery disease; COPD = chronic obstructive pulmonary disease; LHF = left heart failure.

<sup>a</sup> For numerical variables, mean ± SD; for categorical variables, number and percentages.

clinical success was obtained in all cases: open conversion was not required. Mean operation time was 107 ± 63 minutes (range 30–300; median 90 minutes) and the mean aortic coverage was 200 ± 9 mm (range 10–38; median 200 mm). Median blood loss was 200 mL (range 50–2,600 mL) and mean blood transfusion was 4 ± 4 units of packed red blood cells (range 0–27; median, 4). ICU stay was < 2 days in 13 (23%) cases, between 3 and 14 days in 24 (43%), and > 15 days in 19 (34%).

### Hemothorax management

Left HTX was present in all cases and was bilateral in three (5%). Hemothorax thickness on pre-operative computed tomography angiography was < 2 cm in 13 (23%) cases, between 2 and 3 cm in 13 (23%), and > 3 cm in 30 (54%). Overall, HTX was drained in 21 (37.59%) cases: 1 of 13 (7%) for HTX < 2 cm, 6 of 13 (46%) for HTX between 2 and 3 cm, and 14 of 30 (47%) for HTX > 3 cm. All patients but one were drained within the first 3 post-operative days. A chest tube was used in 16 (28%), and surgical evacuation in six (11%). Co-morbidities and risk factors were similar in drained and retained HTX, but drained patients showed worst pre-operative levels of lactate (mEq/L, 3.1 ± 2.3 vs. 1.8 ± 1.0, *p* = .028), PCO<sub>2</sub> (mmHg, 41 ± 9 vs. 36 ± 5, *p* = .037), and post-operative RI (198 ± 106 vs. 311 ± 88, *p* = .001): in hospital mortality did not differ in these two groups (18% vs. 6%, *p* = .197). Although traumatic and non-traumatic ruptures had significantly different co-morbidities, risk factors and respiratory status (Table 2), the need of chest drainage (48% vs. 33%, *p* = .838) did not differ between the two groups.

### Early outcomes

The aortic related mortality rate was 12.5% (*n* = 7): causes of death were shock (*n* = 6, 11%) and respiratory failure (*n* = 1, 2%). Mean hospitalization was 21 ± 16 days (range 2–83 days). TEVAR related complications are summarized in Table 3. In particular, respiratory complications occurred in

**Table 2.** Comparative data of aortic pathology ruptures.

| Variable <sup>a</sup>         | Traumatic<br>( <i>n</i> = 23) | Non-traumatic<br>( <i>n</i> = 33) | <i>p</i> |
|-------------------------------|-------------------------------|-----------------------------------|----------|
| Demographic data (%)          |                               |                                   |          |
| M : F                         | 19 : 4                        | 21 : 12                           | .145     |
| Age (mean ± SD)               | 46 ± 20                       | 74 ± 11                           | < .001   |
| IHD                           | 3 (13)                        | 13 (39)                           | .039     |
| Cardiac disease               | 0 (0)                         | 7 (21)                            | .034     |
| COPD                          | 4 (17)                        | 12 (36)                           | .145     |
| Diabetes                      | 1 (4)                         | 11 (33)                           | .009     |
| AKI                           | 0 (0)                         | 8 (24)                            | .016     |
| Obesity (BMI > 30)            | 1 (4)                         | 13 (39)                           | .004     |
| Operative risk                |                               |                                   |          |
| Shock (%)                     | 13 (56.5)                     | 6 (18)                            | .004     |
| EuroSCORE<br>(mean ± SD)      | 8 ± 2.5                       | 10 ± 4                            | .038     |
| Respiratory parameters (± SD) |                               |                                   |          |
| Acid base status              | 7.3 ± 0.1                     | 7.4 ± 0.1                         | < .001   |
| Po <sub>2</sub> (mmHg)        | 93 ± 23                       | 82 ± 22                           | .076     |
| Pco <sub>2</sub> (mmHg)       | 41 ± 9                        | 37 ± 6                            | .050     |
| Lactates (mEq/L)              | 3.3 ± 2.5                     | 2 ± 1.4                           | .016     |
| Hemothorax thickness (%)      |                               |                                   |          |
| < 2 cm                        | 2 (9)                         | 11 (33)                           | .051     |
| 2–3 cm                        | 3 (13)                        | 11 (33)                           | .119     |
| > 3 cm                        | 18 (78)                       | 11 (33)                           | < .001   |
| Respiratory<br>index (± SD)   | 208 ± 104                     | 286 ± 105                         | .008     |

AKI = acute kidney injury; BMI = body mass index; CAD = coronary artery disease; COPD = chronic obstructive pulmonary disease; Hb = hemoglobin; ICU = intensive care unit; LHF = left heart failure; LOS = length of stay; SBP = systolic blood pressure.

<sup>a</sup> For numerical variables, mean ± SD; for categorical variables, number and percentages.

23 (41%) patients. Overall, no patients required definitive tracheotomy; briefly, those who developed post-operative respiratory complications were older (63 ± 21 vs. 62 ± 21, *p* = .012), had lower pre-operative PO<sub>2</sub> values (mmHg, 80 ± 24 vs. 91 ± 21, *p* = .012), and worse RI at the end of intervention (265 ± 110 vs. 311 ± 88, *p* = .001). No other significant differences were observed (Table 4). Respiratory complications led to longer mechanical ventilation (7 ± 9 vs. 2 ± 7 days, *p* = .001), ICU stay (2.3 ± 0.7 vs.

**Table 3.** Post-operative major complications.

| Complication        | <i>n</i> (%) | Treatment         |
|---------------------|--------------|-------------------|
| Respiratory         |              |                   |
| Grade 1             | 8 (14)       |                   |
| Grade 2             | 8 (14)       |                   |
| Grade 3             | 7 (12.5)     |                   |
| Cardiovascular      |              |                   |
| Atrial fibrillation | 2 (3.5)      |                   |
| Ischemic colitis    | 1 (2)        | Hartman procedure |
| AVF malfunction     | 1 (2)        | vBCT stent        |
| Miscellaneous       |              |                   |
| Paraplegia          | 1 (2)        |                   |
| AKI                 | 1 (2)        | Temporary HD      |
| Dysphagia           | 1 (2)        |                   |

AVF = arteriovenous fistula; AKI = acute kidney injury; vBCT = vein brachiocephalic trunk; HD = hemodialysis.

1.9 ± 0.8 days,  $p = .017$ ), and hospitalization (296 ± 21 vs. 17 ± 12 days,  $p = .009$ ). Post-operatively, traumatic and non-traumatic ruptures did not differ significantly for prolonged (> 10 days) ventilation time (9% vs. 12%,  $p = 1.0$ ), respiratory complications (30% vs. 48%,  $p = .269$ ), hospitalization (25 ± 21 vs. 19 ± 15 days,  $p = .217$ ), and in hospital mortality (9% vs. 12%,  $p = 1.0$ ). In hospital mortality did not differ between patients with and without respiratory complications (13% vs. 12%;  $p = .990$ ).

### Late outcomes

Forty-nine (87.5%) patients were discharged alive: 48 of 49 (86%) patients were available for follow up. Mean follow up was 39 ± 29 months (range 1–108; median 40 months). TEVAR related re-intervention was never required. Nine (19%) patients died during follow up (mean 24 months): aortic related mortality was not observed, and the predominant cause of death was acute myocardial infarction ( $n = 4$ ). Estimated overall survival was 85 ± 4.8% at 3 months, 79 ± 5.6% at 1 year, and 61 ± 8.5% at 5 years.

### DISCUSSION

Population based analysis and single center studies have reported a significant benefit of TEVAR versus open repair for peri-operative mortality, regardless of the indication for repair and type of pathology.<sup>1,2,5–7</sup> Nevertheless, peri-operative mortality after TEVAR for thoracic aortic rupture is still disturbingly high in the range of 8–30%.<sup>1,5,6,22–25</sup> The 12.5% in hospital mortality rate is good if compared with the 35.3% predicted pre-operatively, and it is consistent

with the average 19.4% emerging from previous published series which have included all aortic pathologies as underlying causes of the thoracic aortic rupture.<sup>1,5,6,8,22,23</sup> By contrast, the long-term benefit of TEVAR has been associated with conflicting results. Goodney et al.<sup>21</sup> found that patients selected for TEVAR had worse long-term survival than patients selected for open repair. However, they studied an aneurysm cohort only, and postulated that TEVARs were offered to higher risk patients. The estimated 61% survival at 5 years in the present study is better than the 26% reported by Goodney et al., and compares favorably with other series (in the range 23–55%) which include different aortic catastrophes.<sup>2,5,6,8,22–25</sup> In an emergent setting, the rapidity and minimal invasiveness of TEVAR may be the preferred approach for patients who may not have tolerated a complex and risky operation.

On admission, HTX has frequently been noted after thoracic aortic rupture with an associated increased 30 day mortality.<sup>1,8,9,14,22</sup> Unfortunately, not all of the studies described the treatment and fate of HTX after TEVAR. In their multicenter study, Jonker et al.<sup>8</sup> reported a 41.4% rate of HTX on admission, and a six fold increased risk of in hospital mortality when it was present after rupture of a descending aneurysm; however, the management and outcomes of HTX were not described. Only Shu et al.<sup>9</sup> reported the treatment of HTX after TEVAR. They drained 22% (6/27) of the cases in order to improve respiratory function, but included a cohort of complicated dissections only and almost always used thoracocentesis to evacuate the pleural hematoma. This study is substantially different: the

**Table 4.** Intra-operative and post-operative details with reference to post-operative respiratory complication.

| Variable <sup>a</sup>       | All patients ( $n = 56$ ) | No complication ( $n = 33$ ) | Complicated ( $n = 23$ ) | $p$  |
|-----------------------------|---------------------------|------------------------------|--------------------------|------|
| Demographic data            |                           |                              |                          |      |
| M : F                       | 40 : 16                   | 25 : 8                       | 15 : 8                   | .55  |
| Age (yrs)                   | 63 ± 21                   | 62 ± 21                      | 64 ± 21                  | .012 |
| Risk factors                |                           |                              |                          |      |
| CAD                         | 16 (29)                   | 8 (24)                       | 8 (35)                   | .55  |
| Valve disease/LHF           | 7 (13)                    | 1 (3)                        | 6 (26)                   | .015 |
| COPD                        | 16 (27)                   | 7 (21)                       | 9 (39)                   | .23  |
| Diabetes                    | 13 (23)                   | 8 (24)                       | 5 (22)                   | .99  |
| AKI                         | 8 (14)                    | 5 (15)                       | 3 (13)                   | .99  |
| Obesity (BMI > 30)          | 15 (27)                   | 8 (24)                       | 7 (30)                   | .76  |
| Aortic disease              |                           |                              |                          |      |
| Degenerative                | 33 (59)                   | 17 (52)                      | 16 (70)                  |      |
| Trauma                      | 23 (41)                   | 16 (48)                      | 7 (30)                   |      |
| Operative risk              |                           |                              |                          |      |
| Shock                       | 19 (34)                   | 10 (30)                      | 9 (39)                   | .57  |
| EuroSCORE (additive)        | 9 ± 4                     | 9 ± 3                        | 10 ± 4                   | .44  |
| Preoperative parameters     |                           |                              |                          |      |
| Acid base status            | 7.4 ± 0.1                 | 7.3 ± 0.1                    | 7.4 ± 0.1                | .99  |
| Po <sub>2</sub> (mmHg)      | 87 ± 23                   | 91 ± 21                      | 80 ± 24                  | .012 |
| Pco <sub>2</sub> (mmHg)     | 38 ± 7                    | 36 ± 5                       | 40 ± 9                   | .13  |
| Lactates (mEq/L)            | 2.4 ± 1.9                 | 1.8 ± 1.3                    | 3.1 ± 2.3                | .53  |
| Hemothorax thickness > 3 cm | 29 (52)                   | 20 (61)                      | 9 (39)                   | .17  |
| Respiratory index           | 265 ± 110                 | 311 ± 88                     | 198 ± 106                | .001 |

AKI = acute kidney injury; BMI = body mass index; CAD = coronary artery disease; COPD = chronic obstructive pulmonary disease; Hb = hemoglobin; LHF = left heart failure; SBP = systolic blood pressure.

<sup>a</sup> For numerical variables, mean ± SD; for categorical variables, number and percentages.

percentage of pleural drainage is more than double and either chest tube or open surgery was used. These data may be explained by the fact that only patients with HTX on the baseline CT scan were included, and they underscore the different severity of the HTX in these cases. Nevertheless, this study agrees with that of Shu et al. that prompt drainage should be performed to improve respiratory function and survival. The aggressive approach may have been a contributing factor to explain the fact that the mortality rate of patients who needed drainage of their HTX was not different from those that were not drained, despite a more critical pre-operative respiratory condition. An important aspect to be analyzed is the meaning of HTX and the need for chest drainage in different settings. Eggebrecht et al.<sup>22</sup> showed that patients with complex aortic pathologies were at higher risk of a poor outcome. In this study, rupture of degenerative disease and traumatic rupture were significantly different populations, but the underlying pathology had no impact on post-operative outcomes. These data are supported by the study of Cambria et al.<sup>6</sup> on thoracic aortic catastrophes when outcomes were controlled for baseline co-morbidities and age, the effect of pathology was not significant. This is not because traumatic patients did worse, rather, the explanation lies in the different risk profile of the two groups: most of the traumatic ruptures occurred in younger patients, whereas patients with ruptures of degenerative disease were older and with multiple more severe co-morbidities. Hence, despite the less severe pre-operative condition, the latter group included more fragile patients who benefited from an equally aggressive drainage of HTX.

An interesting issue to be debated is how often there is rupture after acute type B dissection. This is not a rare event, and has been associated with a higher operative mortality than those with other types of complications.<sup>26–29</sup> The 14% in this study is less than the 23.1–60% reported in different types of studies, but it is similar to the 15.2% reported by Cambria et al.,<sup>6</sup> who also included all types of thoracic aortic ruptures.<sup>26–29</sup>

It has been demonstrated that when lysis of chest hematomas is incomplete, by the 7th day HTX may adhere to the lung and pleura, making it difficult to remove, and leading to respiratory complications or infection.<sup>30</sup> Therefore, clotted HTX should be considered for evacuation within a reasonable time; currently, it is still unclear whether, when, and how to perform chest drainage. A recent paper by Bradley et al.<sup>31</sup> showed that the volume of retained HTX was not predictive of post-operative respiratory complications. Shu et al.<sup>9</sup> drained the chest according to the amount of HTX specifically to improve respiratory function and obtained resolution in all cases. These experiences analyzed only specific populations such as non-aorta related trauma patients or dissections, did not use a specific timing algorithm or volume limit for HTX evacuation, or report the respiratory outcomes. All these data underscore the lack of uniformity in current practice for HTX management. This potentially also applies to this study, because 40% of the cases were drained. Nevertheless, the policy was

to drain the chest promptly in the post-operative period, especially for the greater HTX thickness (> 3 cm) and in patients with a respiratory index < 200. Certainly, the compared groups are not similar, but such an aggressive approach in this subset of patients may have contributed to limiting the number of the most severe (SVS grade 2 or 3) respiratory complications and overall mortality. The incidence of respiratory complications after TEVAR for thoracic aortic rupture has been reported in the range of 15–39%.<sup>2,4,5,22,25</sup> A recent nationwide study on use and outcomes of TEVAR for ruptured thoracic aortic aneurysms reported that respiratory complications were alarmingly high at 63%, and even higher in the TEVAR group compared with conventional repair.<sup>3</sup> However, when pleural effusion or HTX were eliminated from the final analysis the respiratory complications were lower in the TEVAR group. None of these studies used a specific definition or analyzed the respiratory complication rate after TEVAR. The 41% reported in this study was higher than previously reported, but all types of respiratory complication should be included in accordance with the SVS guidelines.<sup>20</sup>

### Limitations of the study

The multicenter design of this study has several limitations. The number of patients remains relatively small for adequate analysis and multivariate adjustment. Another important limitation was the retrospective design. Finally, owing to missing worldwide accepted guidelines, indications for the management of the retained HTX may have affected the results. However, due to the rarity of these events and its emergent nature, it will be very difficult to ever present a large and homogeneous study. Despite these limitations, there are no similar studies in the literature: the major critical aspect of the paper is the inability to present generalizable conclusions, but this cohort of patients with HTX due to thoracic aortic rupture treated with TEVAR is currently the largest available, and may lead to larger studies to outline the best management of this condition.

### CONCLUSION

This study confirms that descending thoracic aortic rupture still carries a high mortality even in the TEVAR era; in addition, respiratory complications have not been eliminated by this less traumatic technique.

Prompt HTX evacuation may have contributed to limiting post-operative mortality in drained patients who presented with significantly worse pre-operative respiratory parameters. Although traumatic and degenerative ruptures are two significantly different scenarios, survival and respiratory outcomes were not affected by the underlying aortic disease.

### CONFLICT OF INTEREST

None.

### FUNDING

None.

## REFERENCES

- 1 Patel HJ, Williams DM, Upchurch Jr GR, Dasika NL, Deeb GM. A comparative analysis of open and endovascular repair for the ruptured descending thoracic aorta. *J Vasc Surg* 2009;**50**(6):1265–70.
- 2 Conrad MF, Ergul EA, Patel VI, Paruchuri V, Kwolek CJ, Cambria RP. Management of diseases of the descending thoracic aorta in the endovascular era: a Medicare population study. *Ann Surg* 2010;**252**(4):603–10.
- 3 Jonker FH, Verhagen HJ, Lin PH, Heijmen RH, Trimarchi S, Lee WA, et al. Open surgery versus endovascular repair of ruptured thoracic aortic aneurysms. *J Vasc Surg* 2011;**53**(5):1210–6.
- 4 Gopaldas RR, Dao TK, LeMaire SA, Huh J, Coselli JS. Endovascular versus open repair of ruptured descending thoracic aortic aneurysms: a nationwide risk-adjusted study of 923 patients. *J Thorac Cardiovasc Surg* 2011;**142**(5):1010–8.
- 5 Naughton PA, Park MS, Morasch MD, Rodriguez HE, Garcia-Toca M, Wang CE, et al. Emergent repair of acute thoracic aortic catastrophes: a comparative analysis. *Arch Surg* 2012;**147**(3):243–9.
- 6 Cambria RP, Crawford RS, Cho JS, Bavaria J, Farber M, Lee WA, et al. A multicenter clinical trial of endovascular stent graft repair of acute catastrophes of the descending thoracic aorta. *J Vasc Surg* 2009;**50**(6):1255–64.
- 7 Murad MH, Rizvi AZ, Malgor R, Carey J, Alkatib AA, Erwin PJ, et al. Comparative effectiveness of the treatments for thoracic aortic transection. *J Vasc Surg* 2011;**53**(1):193–9.
- 8 Jonker FH, Verhagen HJ, Lin PH, Heijmen RH, Trimarchi S, Lee WA, et al. Outcomes of endovascular repair of ruptured descending thoracic aortic aneurysms. *Circulation* 2010;**121**(25):2718–23.
- 9 Shu C, He H, Li QM, Li M, Jiang XH, Luo MY. Endovascular repair of complicated acute type-B aortic dissection with stentgraft: early and mid-term results. *Eur J Vasc Endovasc Surg* 2011;**42**(4):448–53.
- 10 Tokui T, Shimono T, Kato N, Hirano T, Takeda K, Yada I. Less invasive therapy using endovascular stent graft repair and video-assisted thoracoscopic surgery for ruptured acute aortic dissection. *Jpn J Thorac Cardiovasc Surg* 2000;**48**(9):603–6.
- 11 Vassiliu P, Velmahos GC, Toutouzias KG. Timing, safety, and efficacy of thoracoscopic evacuation of undrained post-traumatic hemothorax. *Am Surg* 2001;**67**(12):1165–9.
- 12 Monaco M, Mondello B, Spinelli F, Gaeta R, Carella G, Lentini S, et al. Endovascular repair of aortic lesions associated to video-assisted thoracoscopy surgery after chest trauma. *Minerva Chir* 2008;**63**(5):329–34.
- 13 Vollroth M, Seeburger J, Kiefer P, Hoebartner M, Bausback Y, Garbade J, et al. Interdisciplinary team approach for complicated type B aortic dissection with concomitant hemothorax by endovascular stent grafting and left side mini thoracotomy: a case report. *J Cardiothorac Surg* 2012;**7**:111.
- 14 Trimarchi S, Segreti S, Grassi V, Lomazzi C, de Vincentiis C, Rampoldi V. Emergent treatment of aortic rupture in acute type B dissection. *Ann Cardiothorac Surg* 2014;**3**(3):319–24.
- 15 Sganga G, Siegel JH, Coleman B, Giovannini I, Boldrini G, Pittiruti M. The physiologic meaning of the respiratory index in various types of critical illness. *Circ Shock* 1985;**17**(3):179–93.
- 16 Clarck RE. Committee to develop a national database for thoracic surgeons. Definitions of terms of the Society of Thoracic Surgeons national cardiac surgery database. *Ann Thor Surg* 1994;**58**:271–3.
- 17 Marone EM, Kahlberg A, Tshomba Y, Chiesa R. Single center experience with endovascular treatment of acute blunt thoracic aortic injuries. *J Cardiovasc Surg* 2013;**54**(1):123–31.
- 18 Lee WA, Matsumura JS, Mitchell RS, Farber MA, Greenberg RK, Azzizadeh A, et al. Endovascular repair of traumatic thoracic aortic injury: clinical practice guidelines of the Society for Vascular Surgery. *J Vasc Surg* 2011;**53**(1):187–92.
- 19 Roques F, Nashef SA, Michel P, Gauducheau E, de Vincentiis C, Baudet E, et al. Risk factors and outcome in European cardiac surgery: analysis of the EuroSCORE multinational database of 19030 patients. *Eur J Cardiothorac Surg* 1999;**15**(6):816–22.
- 20 Fillinger MF, Greenberg RK, McKinsey JF, Chaikof EL. Society for Vascular Surgery Ad Hoc Committee on TEVAR Reporting Standards. Reporting standards for thoracic endovascular aortic repair (TEVAR). *J Vasc Surg* 2010;**52**(4):1022–33.
- 21 Goodney PP, Travis L, Lucas FL, Fillinger MF, Goodman DC, Cronenwett JL, et al. Survival after open versus endovascular thoracic aortic aneurysm repair in an observational study of the Medicare population. *Circulation* 2011;**124**(24):2661–9.
- 22 Eggebrecht H, Schmermund A, Herold U, Baumgart D, Martini S, Kuhnt O, et al. Endovascular stent graft placement for acute and contained rupture of the descending thoracic aorta. *Catheter Cardiovasc Interv* 2005;**66**(4):474–8.
- 23 Alsac JM, Pierard T, El Batti S, Acouch P, Julia P, Fabiani JN. Applicability and mid-term results of endovascular treatment for descending thoracic acute aortic syndromes. *Ann Vasc Surg* 2013;**27**(8):1029–35.
- 24 Mitchell ME, Rushton Jr FW, Boland AB, Byrd TC, Baldwin ZK. Emergency procedures on the descending thoracic aorta in the endovascular era. *J Vasc Surg* 2011;**54**(5):1298–302.
- 25 Geisbüsch P, Kotelis D, Weber TF, Hyhlik-Dürr A, Böckler D. Endovascular repair of ruptured thoracic aortic aneurysms is associated with high peri-operative mortality and morbidity. *J Vasc Surg* 2010;**51**(2):299–304.
- 26 Trimarchi S, Nienaber CA, Rampoldi V, Myrmet T, Suzuki T, Bossone E, et al. IRAD Investigators. Role and results of surgery in acute type B aortic dissection: insights from the International Registry of Acute Aortic Dissection (IRAD). *Circulation* 2006;**114**(1 Suppl):1357–64.
- 27 Steuer J, Eriksson MO, Nyman R, Björck M, Wanhainen A. Early and long-term outcome after thoracic endovascular aortic repair (TEVAR) for acute complicated type B aortic dissection. *Eur J Vasc Endovasc Surg* 2011;**41**(3):318–23.
- 28 White RA, Miller DC, Criado FJ, Dake MD, Diethrich EB, Greenberg RK, et al. Multidisciplinary Society for Vascular Surgery Outcomes Committee. Report on the results of thoracic endovascular aortic repair for acute, complicated, type B aortic dissection at 30 days and 1 year from a multidisciplinary subcommittee of the Society for Vascular Surgery Outcomes Committee. *J Vasc Surg* 2011;**53**(4):1082–90.
- 29 Wiedemann D, Ehrlich M, Amabile P, Lovato L, Rousseau H, Evangelista-Masip A, et al. Emergency endovascular stent grafting in acute complicated type B dissection. *J Vasc Surg* 2014;**60**(5):1204–8.
- 30 Boersma WG, Stigt JA, Smit HJ. Treatment of haemothorax. *Respir Med* 2010;**104**(11):1583–7.
- 31 Bradley M, Okoye O, DuBose J, Inaba K, Demetriades D, Scalea T, et al. Risk factors for post-traumatic pneumonia in patients with retained haemothorax: results of a prospective, observational AAST study. *Injury* 2013;**44**(9):1159–64.